



LAB #: F000000-0000-0
 PATIENT: Sample Patient
 ID: P00000000
 SEX: Female
 AGE: 50

CLIENT #: 12345
 DOCTOR:
 Doctor's Data, Inc.
 3755 Illinois Ave.
 St. Charles, IL 60174

Comprehensive Parasitology, stool, x3

BACTERIOLOGY CULTURE

Expected/Beneficial flora

1+ Bacteroides fragilis group
 4+ Bifidobacterium spp.
 NG Escherichia coli
 1+ Lactobacillus spp.
 NG Enterococcus spp.

 2+ Clostridium spp.
 NG = No Growth

Commensal (Imbalanced) flora

1+ Gamma hemolytic strep
 3+ Hemolytic Escherichia coli

Dysbiotic flora

BACTERIA INFORMATION

Expected /Beneficial bacteria make up a significant portion of the total microflora in a healthy & balanced GI tract. These beneficial bacteria have many health-protecting effects in the GI tract including manufacturing vitamins, fermenting fibers, digesting proteins and carbohydrates, and propagating anti-tumor and anti-inflammatory factors.

Clostridia are prevalent flora in a healthy intestine. Clostridium spp. should be considered in the context of balance with other expected/beneficial flora. Absence of clostridia or over abundance relative to other expected/beneficial flora indicates bacterial imbalance. If *C. difficile* associated disease is suspected, a Comprehensive Clostridium culture or toxigenic *C. difficile* DNA test is recommended.

Commensal (Imbalanced) bacteria are usually neither pathogenic nor beneficial to the host GI tract. Imbalances can occur when there are insufficient levels of beneficial bacteria and increased levels of commensal bacteria. Certain commensal bacteria are reported as dysbiotic at higher levels.

Dysbiotic bacteria consist of known pathogenic bacteria and those that have the potential to cause disease in the GI tract. They can be present due to a number of factors including: consumption of contaminated water or food, exposure to chemicals that are toxic to beneficial bacteria; the use of antibiotics, oral contraceptives or other medications; poor fiber intake and high stress levels.

YEAST CULTURE

Normal flora

No yeast isolated

Dysbiotic flora

MICROSCOPIC YEAST

Result:

Rare

Expected:

None - Rare

The microscopic finding of yeast in the stool is helpful in identifying whether there is proliferation of yeast. Rare yeast may be normal; however, yeast observed in higher amounts (few, moderate, or many) is abnormal.

YEAST INFORMATION

Yeast normally can be found in small quantities in the skin, mouth, intestine and mucocutaneous junctions. Overgrowth of yeast can infect virtually every organ system, leading to an extensive array of clinical manifestations. Fungal diarrhea is associated with broad-spectrum antibiotics or alterations of the patient's immune status. Symptoms may include abdominal pain, cramping and irritation. When investigating the presence of yeast, disparity may exist between culturing and microscopic examination. Yeast are not uniformly dispersed throughout the stool, this may lead to undetectable or low levels of yeast identified by microscopy, despite a cultured amount of yeast. Conversely, microscopic examination may reveal a significant amount of yeast present, but no yeast cultured. Yeast does not always survive transit through the intestines rendering it unviable.

Comments:

Date Collected: 11/30/2011
 Date Received: 12/1/2011
 Date Completed: 12/9/2011

* *Aeromonas, Campylobacter, Plesiomonas, Salmonella, Shigella, Vibrio, Yersinia, & Edwardsiella tarda* have been specifically tested for and found absent unless reported.

v5.09



LAB #: F000000-0000-0
 PATIENT: Sample Patient
 ID: P000000-0000-0
 SEX: Female
 AGE: 50

CLIENT #: 12345
 DOCTOR:
 Doctor's Data, Inc.
 3755 Illinois Ave.
 St. Charles, IL 60174

Comprehensive Parasitology, stool, x3

PARASITOLOGY/MICROSCOPY *

Sample 1

Mod Blastocystis hominis

Sample 2

Mod Blastocystis hominis

Sample 3

Few Blastocystis hominis
 Rare Yeast

*A trichrome stain and concentrated iodine wet mount slide is read for each sample submitted.

PARASITOLOGY INFORMATION

Intestinal parasites are abnormal inhabitants of the gastrointestinal tract that have the potential to cause damage to their host. The presence of any parasite within the intestine generally confirms that the patient has acquired the organism through fecal-oral contamination. Factors such as contaminated food and water supplies, day care centers, increased international travel, pets, carriers such as mosquitoes and fleas, and sexual transmission have contributed to an increased prevalence of intestinal parasites. It is estimated that close to one billion people worldwide are infected. Damage to the host includes parasitic burden, migration, blockage and pressure. Immunologic inflammation, hypersensitivity reactions and cytotoxicity also play a large role in the morbidity of these diseases. The infective dose often relates to severity of the disease and repeat encounters can be additive.

There are two main classes of intestinal parasites that can cause human intestinal disease. They include protozoa and helminths. The protozoa typically have two stages; the trophozoite stage that is the metabolically active, invasive stage and the cyst stage, which is the vegetative inactive form resistant to unfavorable environmental conditions outside the human host. Helminths are large, multicellular organisms that are generally visible to the naked eye in their adult stages. Like protozoa, helminths can be either free-living or parasitic in nature. In their adult form, helminths cannot multiply in humans.

In general, acute manifestations of parasitic infection may involve diarrhea with or without mucus and or blood, fever, nausea, or abdominal pain. However these symptoms do not always occur. Consequently, parasitic infections may not be diagnosed or eradicated. If left untreated, chronic parasitic infections can cause damage to the intestinal lining and can be an unsuspected cause of illness and fatigue. Chronic parasitic infections can also be associated with increased intestinal permeability, irritable bowel syndrome, irregular bowel movements, malabsorption, gastritis or indigestion, skin disorders, joint pain, allergic reactions, and decreased immune function.

In some instances, parasites may enter the circulation and travel to various organs causing severe organ diseases such as liver abscesses and cysticercosis. In addition, some larval migration can cause pneumonia and in rare cases hyper infection syndrome with large numbers of larvae being produced and found in every tissue of the body.

GIARDIA/CRYPTOSPORIDIUM IMMUNOASSAY

	Within	Outside	Reference Range
Giardia lamblia	Neg	Neg	Neg
Cryptosporidium	Neg	Neg	Neg

Giardia lamblia is flagellated protozoan that infects the small intestine and is passed in stool and spread by the fecal-oral route. Waterborne transmission is the major source of giardiasis.

Cryptosporidium is a coccidian protozoa that can be spread from direct person-to-person contact or waterborne transmission.

Comments:

Date Collected: 11/30/2011
 Date Received: 12/1/2011
 Date Completed: 12/9/2011

v5.09

INTRODUCTION

This analysis of the stool specimen provides fundamental information about the overall gastrointestinal health of the patient. When abnormal microflora or significant aberrations in intestinal health markers are detected, specific interpretive paragraphs are presented. If no significant abnormalities are found, interpretive paragraphs are not presented.

Beneficial Flora

One or more of the expected (beneficial) bacteria are low in this specimen. Beneficial flora include lactobacilli, bifidobacteria, clostridia, *Bacteroides fragilis* group, enterococci, and some strains of *Escherichia coli*. The beneficial flora have many health-protecting effects in the gut, and as a consequence, are crucial to the health of the whole organism. Some of the roles of the beneficial flora include digestion of proteins and carbohydrates, manufacture of vitamins and essential fatty acids, increase in the number of immune system cells, break down of bacterial toxins and the conversion of flavinoids into anti-tumor and anti-inflammatory factors. Lactobacilli, bifidobacteria, clostridia, and enterococci secrete lactic acid as well as other acids including acetate, propionate, butyrate, and valerate. This secretion causes a subsequent decrease in intestinal pH, which is crucial in preventing an enteric proliferation of microbial pathogens, including bacteria and yeast. Many GI pathogens thrive in alkaline environments. Lactobacilli also secrete the antifungal and antimicrobial agents lactocidin, lactobacillin, acidolin, and hydrogen peroxide. The beneficial flora of the GI have thus been found useful in the inhibition of microbial pathogens, prevention and treatment of antibiotic associated diarrhea, prevention of traveler's diarrhea, enhancement of immune function, and inhibition of the proliferation of yeast.

In a healthy balanced state of intestinal flora, the beneficial flora make up a significant proportion of the total microflora. Healthy levels of each of the beneficial bacteria are indicated by either a 3+ or 4+ (0 to 4 scale). However, some individuals have low levels of beneficial bacteria and an overgrowth of nonbeneficial (imbalances) or even pathogenic microorganisms (dysbiosis). Often attributed to the use of antibiotics, individuals with low beneficial bacteria may present with chronic symptoms such as irregular transit time, irritable bowel syndrome, bloating, gas, chronic fatigue, headaches, autoimmune diseases (e.g., rheumatoid arthritis), and sensitivities to a variety of foods. Treatment may include the use of probiotic supplements containing various strains of lactobacilli, bifidobacteria and enterococci and consumption of cultured or fermented foods including yogurt, kefir, miso, tempeh and tamari sauce. Polyphenols in green and ginseng tea have been found to increase the numbers of beneficial bacteria. If dysbiosis is present, treatment may also include the removal of pathogenic bacteria, yeast, or parasites.

Percival M. Intestinal Health. *Clin Nutr In.* 1997;5(5):1-6.

Fuller R. Probiotics in Human Medicine. *Gut.* 1991;32: 439-442.

Siitonen S, Vapaatalo H, Salminen S, et al. Effect of Lactobacilli GG Yoghurt in Prevention of Antibiotic Associated Diarrhea. *Ann Med.* 1990; 22:57-59.

Oksanen P, Salminen S, Saxelin M, et al. Prevention of Travelers' Diarrhea by Lactobacillus GG. *Ann Med.* 1990; 22:53-56.

Perdigon G, Alvarez M, et al. The Oral Administration of Lactic Acid Bacteria Increases the Mucosal Intestinal Immunity in Response to Enteropathogens. *J Food Prot.* 1990;53:404-410.

Valeur, N, et al. Colonization and Immunomodulation by Lactobacillus reuteri ATCC 55730 in the Human Gastrointestinal Tract. *Appl Environ. Microbiol.* 2004 Feb; 70(2):1176-81.

Elmer G, Surawicz C, and McFarland L. Biotherapeutic agents - a Neglected Modality for the Treatment and Prevention of Intestinal and Vaginal Infections. *JAMA.* 1996; 275(11):870-876.

Fitzsimmons N and Berry D. Inhibition of Candida albicans by Lactobacillus acidophilus: Evidence for Involvement of a Peroxidase System. *Microbio.* 1994; 80:125-133

Weisburger JH. *Proc Soc Exp Biol Med* 1999;220(4):271-5.

Imbalanced flora

Imbalanced flora are those bacteria that reside in the host gastrointestinal tract and neither injure nor benefit the host. Certain dysbiotic bacteria may appear under the imbalances category if found at low levels because they are not likely pathogenic at the levels detected. When imbalanced flora appear, it is not uncommon to find inadequate levels of one or more of the beneficial bacteria and/or a fecal pH which is more towards the alkaline end of the reference range (6.5 - 7.2). It is also not uncommon to find hemolytic or mucoid E. coli with a concomitant deficiency of beneficial E. coli and alkaline pH, secondary to a mutation of beneficial E. coli in alkaline conditions (DDI observations). Treatment with antimicrobial agents is unnecessary unless bacteria appear under the dysbiotic category.

Mackowiak PA. The normal microbial flora. *N Engl J Med.* 1982;307(2):83-93.

Microscopic yeast

Microscopic examination has revealed yeast in this stool sample. The microscopic finding of yeast in the stool is helpful in identifying whether the proliferation of fungi, such as Candida albicans, is present. Yeast is normally found in very small amounts in a healthy intestinal tract. While small quantities of yeast (reported as none or rare) may be normal, yeast observed in higher amounts (few, moderate to many) is considered abnormal.

An overgrowth of intestinal yeast is prohibited by beneficial flora, intestinal immune defense (secretory IgA), and intestinal pH. Beneficial bacteria, such as Lactobacillus colonize in the intestines and create an environment unsuitable for yeast by producing acids, such as lactic acid, which lowers intestinal pH. Also, lactobacillus is capable of releasing antagonistic substances such as hydrogen peroxide, lactocidin, lactobacillin, and acidolin.

Many factors can lead to an overgrowth of yeast including frequent use of antibiotics (leading to insufficient beneficial bacteria), synthetic corticosteroids, oral contraceptives, and diets high in sugar. Although there is a wide range of symptoms which can result from intestinal yeast overgrowth, some of the most common include brain fog, fatigue, recurring vaginal or bladder infections, sensitivity to smells (perfumes, chemicals, environment), mood swings/depression, sugar and carbohydrate cravings, gas/bloating, and constipation or loose stools.

A positive yeast culture (mycology) and sensitivity to prescriptive and natural agents is helpful in determining which anti-fungal agents to use as part of a therapeutic treatment plan for chronic colonic yeast. However, yeast are colonizers and do not appear to be dispersed uniformly throughout the stool. Yeast may therefore be observed microscopically, but not grow out on culture even when collected from the same bowel movement.

Parasites

Parasites were detected by microscopic examination in this stool specimen. Intestinal parasites are abnormal inhabitants of the GI tract that live off and have the potential to cause damage to their host. Factors such as contaminated food and water supplies, day care centers, increased international travel, pets, carriers such as mosquitoes and fleas, and sexual transmission have contributed to an increased prevalence of intestinal parasites.

In general, acute manifestations of parasitic infection may involve diarrhea with or without mucus and/or blood, fever, nausea, or abdominal pain. However, these symptoms do not always occur. Consequently, parasitic infections may not be diagnosed and eradicated. If left untreated, chronic parasitic infections can cause damage to the intestinal lining and can be an unsuspected cause of illness and fatigue. Chronic parasitic infections can also be associated with increased intestinal permeability, irritable bowel syndrome, irregular bowel movements, malabsorption, gastritis or indigestion, skin disorders, joint pain, allergic reactions, decreased immune function, and fatigue.

Murray MT. Stomach Ailments And Digestive Disturbances. Rocklin, CA: Prima Publishing;1997.

Gittleman AL. Guess What Came to Dinner Parasites And Your Health. New York, NY: Penguin Group; 2001.

Blastocystis hominis

Blastocystis hominis was identified in this specimen. Blastocystis hominis is a common protozoan found throughout the world. Blastocystis is transmitted via the fecal-oral route or from contaminated food or water.

Whether Blastocystis infection can cause symptoms is still considered controversial. Symptoms may be compounded by concomitant infection with other parasitic organisms, bacteria, or viruses. Often, B. hominis is found along with other such organisms. Nausea, diarrhea, abdominal pain, anal itching, weight loss, and excess gas have been reported in some persons with Blastocystis infection.

Metronidazole has been the traditionally considered the most effective drug (recommended

adult dosage varies from 250 mg bid for 5-7 days to 750 mg tid x 10 days). Iodoquinol is also an effective medication (650 mg tid x 20 days). Recommended therapy can also eliminate *G. lamblia*, *E. histolytica* and *D. fragilis*, all of which may be concomitant undetected pathogens and part of patient symptomatology. Various herbs may be effective, including oil of oregano. Limit refined carbohydrates in diet.

For more information:

1. Albrecht H, Stellbrink HJ, Koperski K, et al. Blastocystis hominis in human immunodeficiency virus-related diarrhea. *Scand J Gastroenterol* 1995;30:909-14.
2. Markell EK, Udkow MP. Blastocystis hominis: pathogen or fellow traveler *Am J Trop Med Hyg* 1986;35:1023-6.
3. Miller RA, Minshew BH. Blastocystis hominis: An organism in search of a disease. *Rev Infect Dis* 1988;10:930-8.
4. Udkow MP, Markell EK. Blastocystis hominis: prevalence in asymptomatic versus symptomatic hosts. *J Infect Dis* 1993;168:242-4.
5. Zuckerman MJ, Watts MT, Ho H., et al. Blastocystis hominis infection and intestinal injury. *Am J Med Sci* 1994;308:96-101.

References:

Sanford JP. *The Sanford Guide to Antimicrobial Therapy*. 35th edition. Gilbert DN, Moellering Jr, RC, Sande MA, eds. Hyde Park (VT): Antimicrobial Therapy Inc; 2005.

Abramowicz, M. *The Medical Letter On Drugs and Therapeutics*. Drugs For Parasitic Infections. New Rochelle (NY): The Medical Letter, Inc.

Beers, M. H., & Berkow, R. (Eds.). *The Merck Manual of Diagnosis and Therapy Online*. <http://www.merck.com/mrkshared/mmanual/section13/chapter161/161a.jsp>, Accessed August, 2005.

CDC Division of Parasitic Diseases website. <http://www.cdc.gov/ncidod/dpd/default.htm>, Accessed August, 2005.

Garcia, LS. *Diagnostic Medical Parasitology*. 4th ed. Washington DC: ASM; 2001; 6.

Leber AL, Movak SM In: Murray PR, Baron EJ, Pfaller MA, Tenover FC, Tenover RH, eds. *Manual of Clinical Microbiology*. 7th ed. Washington DC: ASM Press; 1999; 1401.